

Maryland HIV/AIDS Quarterly Update

Second Quarter 2016

Data reported through June 30, 2016



Center for HIV Surveillance, Epidemiology and Evaluation
Infectious Disease Epidemiology and Outbreak Response Bureau
Prevention and Health Promotion Administration
Maryland Department of Health and Mental Hygiene
<http://phpa.dhmh.maryland.gov/oideor/chse>
1-800-358-9001



TABLE OF CONTENTS

Section I – Background Information	1
HIV/AIDS Reporting Requirements	1
For Assistance with HIV/AIDS Reporting.....	2
Limitations in the HIV/AIDS Data.....	2
Stages of a Case of HIV/AIDS.....	3
Changes in Case Terminology	3
Laboratory Data.....	4
Sources of Data.....	4
Tabulation of Column Totals.....	4
Data Suppression.....	4
Glossary of Terms	4
Section II – Adult/Adolescent Cases by Jurisdiction	6
Table 1 – Adult/Adolescent HIV Diagnoses during 1/1/2015-12/31/2015, First CD4 Test Result, Percent Linked to Care, and Percent Late Diagnosis, by Jurisdiction, Reported through 6/30/2016	6
Table 2 – Adult/Adolescent AIDS Diagnoses during 1/1/2015-12/31/2015, Mean Years from HIV Diagnosis and Percent Late HIV Diagnosis, by Jurisdiction, Reported through 6/30/2016	7
Table 3 – Adult/Adolescent HIV Cases Alive on 12/31/2015, by Jurisdiction, Reported through 6/30/2016.....	8
Table 4 – CD4 Test Results for Adult/Adolescent HIV Cases Alive on 12/31/2015, Reported through 6/30/2016	9
Table 5 – Viral Load Test Results for Adult/Adolescent HIV Cases Alive on 12/31/2015, by Jurisdiction, Reported through 6/30/2016..	10

Section I – Background Information

HIV/AIDS Reporting Requirements

The Maryland HIV/AIDS Reporting Act of 2007 went into effect on April 24, 2007. The law expanded HIV/AIDS reporting and required that HIV cases be reported by name. The following highlights the reporting requirements of Health-General Articles 18-201.1, 18-202.1, and 18-205 of the Annotated Code of Maryland, as specified in COMAR 10.18.02.

- Physicians are required to report patients in their care with diagnoses of HIV or AIDS immediately to the Local Health Department where the physician’s office is located by mailing DHMH Form 1140. Reports are also accepted by phone.
- Physicians are required to report infants born to HIV positive mothers within 48 hours to the State Health Department by mailing DHMH Form 1140. Reports are also accepted by phone.
- Clinical and infection control practitioners in hospitals, nursing homes, hospice facilities, medical clinics in correctional facilities, inpatient psychiatric facilities, and inpatient drug rehabilitation facilities are required to report patients in the care of the institution with diagnoses of HIV or AIDS within 48 hours to the Local Health Department where the institution is located by mailing DHMH Form 1140. Reports are also accepted by phone.

Facilities with large volumes are encouraged to contact the State Health Department to establish electronic reporting.

- Laboratory directors are required to report patients with laboratory results indicating HIV infection (e.g., positive confirmatory HIV diagnostic tests, all CD4 immunological tests, all HIV viral load tests, and all HIV genotype and phenotype tests) within 48 hours to the Local Health Department where the laboratory is located, or if out of state to the Maryland State Health Department, by mailing DHMH Form 4492. Laboratories are encouraged to contact the State Health Department to establish electronic reporting.

Reporting forms and instructions are available on our website:

<http://phpa.dhmh.maryland.gov/oideor/chse/sitepages/reporting-material.aspx>

For Assistance with HIV/AIDS Reporting

For assistance with reporting, including establishment of routine, electronic, or other alternate methods of reporting to the Health Department, please contact the Center for HIV Surveillance, Epidemiology and Evaluation in the Maryland Department of Health and Mental Hygiene at 410-767-5227.

Limitations in the HIV/AIDS Data

This epidemiological profile only contains data for HIV and AIDS cases that have been diagnosed by a health care provider, were reported to the health department by name, and were residents of Maryland at the time of diagnosis. Nationally, it has been estimated that 12.8% of people living with HIV infection are undiagnosed. In Maryland, it is estimated that 18.7% of people living with HIV infection are undiagnosed. In addition, despite a massive effort during which over 17,000 HIV cases were reported after the Maryland HIV reporting law changed on April 24, 2007, not all diagnosed HIV cases previously reported by Maryland's code-based identifier were located and re-reported by name. In addition, many of the re-reported HIV cases were identified by a recent diagnosis and not by their earliest diagnosis, resulting in an under-reporting of HIV diagnoses before 2001 and an over-reporting of HIV diagnoses from 2001 to 2008. Caution should be exercised in using the number of living HIV cases without AIDS and in interpreting trends in the number of reported HIV diagnoses. Furthermore, laboratory data are only available for cases receiving medical care, usually only at facilities in Maryland, and only includes test results that have been reported to the health department.

Please note that not all data has been geocoded in the quarterly reports and therefore is preliminary. Geocoding is the process of assigning geographic identifiers to map features and data records. Addresses are standard data elements required by law and submitted as part of reporting requirements; however, the information may be incomplete which then requires a geocoding process to improve the quality of data. This process is fully completed on the end-of-the-year data set.

Stages of a Case of HIV/AIDS

Untreated HIV disease progresses from HIV infection to AIDS to death. These are biological events that occur whether or not a person receives any medical care. For example, a person can be HIV infected but never have an HIV test and so they do not have an HIV diagnosis. A medical provider diagnoses that these biological events have occurred and records them as a medical event. The law requires medical providers to report these medical events to the Health Department, thereby creating a surveillance event.

Time Point	Biological Event	Medical Event	Surveillance Event
1	HIV Infection		
2		HIV Diagnosis	
3			HIV Report
4	AIDS Conditions		
5		AIDS Diagnosis	
6			AIDS Report
7	Death		
8		Death Diagnosis	
9			Death Report

For surveillance purposes, a case of HIV/AIDS can only move through time in one direction, from HIV infection to death report [from time point 1 to time point 9], but may skip over individual stages. Events can occur simultaneously, but usually there is a time lag between them. The time lag between events can be measured in days, months, or years.

For example, the time between HIV infection [time point 1] and the test that diagnoses HIV [time point 2] may be several years, and it may then take several days for the laboratory and physician to report the diagnosis to the health department [time point 3]. In a second example, a person with diagnosed and reported HIV infection [time point 3] may die [time point 7] without developing AIDS, thereby skipping the three AIDS events (conditions, diagnosis, and report [time points 4, 5 and 6]). And in a third example, a person with undiagnosed HIV infection [time point 1] may become sick, enter the hospital, and die [time point 7] of what is later determined to be AIDS. In that situation, HIV diagnosis [time point 2], AIDS diagnosis [time point 5], and death diagnosis [time point 8] would all occur at the same time, and that would have been many years after the initial HIV infection [time point 1].

Changes in Case Terminology

The terminology for HIV and AIDS cases was changed from earlier epidemiological profiles to be more precise, with Reported Diagnoses replacing Incidence and Living Cases replacing Prevalence. Incidence is a measure of the number of new events (such as HIV infections) in a population during a period of time. Prevalence is a measure of the number of people living with a condition (such as HIV) in a population at a certain time. Prevalence includes both new and old diagnosed cases as well as undiagnosed infections. For HIV, Incidence and Prevalence cannot be directly measured and must be estimated using statistical methods. The HIV surveillance system is able to provide the actual number of diagnoses and deaths that are reported in the population.

For this epidemiological profile, the reports received through a certain time (a quarter-year) are used to generate the number of diagnoses during the prior years. This six-month lag allows for delays in reporting and time to complete investigations. For example, the Reported HIV Diagnoses for 1/1/2015-12/31/2015 are the total of the reported HIV cases with or without an AIDS diagnosis, diagnosed with HIV during 1/1/2015-12/31/2015, as reported by name through 6/30/2016.

To calculate the number of Living Cases we count up all of the Reported Diagnoses from the beginning of the epidemic (all the Reported HIV Diagnoses each year) and subtract all of the Reported Deaths. For example, the Total Living HIV Cases on 12/31/2015 are the total of the reported HIV Cases with or without an AIDS diagnosis and not reported to have died as of 12/31/2015 as reported by name through 6/30/2016.

Laboratory Data

CD4+ T-lymphocyte tests are measures of a person's immune system function. An HIV infected adult is considered to have AIDS if they have less than 200 CD4 cells per microliter of blood. Viral load (VL) tests are measures of the amount of HIV in a person's body. The goal of HIV treatment is to have a very low number of copies of virus per milliliter of blood, below what the test can measure, which is called an undetectable level. Treatment recommendations are that a person in HIV medical care should have their CD4 and VL levels measured regularly, at least once per year. We use the presence of these lab tests as an indicator that someone has been "linked to care" initially after diagnosis or in following years that they remain "in care".

Sources of Data

Information on HIV and AIDS diagnoses, including residence at diagnosis, age, race/ethnicity, sex at birth, country of birth, vital status, HIV exposure category, and CD4 and HIV viral load test results are from the Maryland Department of Health and Mental Hygiene's Enhanced HIV/AIDS Reporting System (eHARS), June 30, 2016.

Population data by sex, age, and race/ethnicity are from the July 1, 2014 U.S. Census Estimates. Due to estimation limitations, some population totals may not equal the sum of its components.

Tabulation of Column Totals

Figures in tables and generally in the text have been rounded. Discrepancies in tables between totals and sums of components are due to rounding.

Data Suppression

In order to protect the confidentiality of reported HIV cases, data are suppressed in the following instances:

- Data describing a demographic group or geographic area (e.g. ZIP code) with a population less than 1,000 people.
- All clinical/laboratory information if it is describing less than 5 cases.
- All exposure/risk information if it is describing less than 5 cases, except in the case of "other" exposure.
- If any cell is suppressed, additional cells are also suppressed as necessary to prevent back calculation of the suppressed cell(s).

Glossary of Terms

Adult/Adolescent Living HIV Cases with AIDS: Reported HIV diagnoses, age 13 years or older at HIV diagnosis, with an AIDS diagnosis, and not reported to have died as of 12/31/2015.

Adult/Adolescent Living HIV Cases without AIDS: Reported HIV diagnoses, age 13 years or older at HIV diagnosis, without an AIDS diagnosis, and not reported to have died as of 12/31/2015.

Adult/Adolescent Reported AIDS Diagnoses: Reported HIV diagnoses, age 13 years or older at HIV diagnosis, with an initial AIDS diagnosis during the specified year.

Adult/Adolescent Reported HIV Diagnoses: Reported HIV diagnoses, age 13 years or older at HIV diagnosis, with an initial HIV diagnosis during the specified year.

Adult/Adolescent Total Living HIV Cases: Reported HIV diagnoses, age 13 years or older at HIV diagnosis, with or without an AIDS diagnosis, and not reported to have died as of 12/31/2015.

CD4 Result Distribution (<200, 200-349, 350-499, 500+): Percent of cases with a CD4 test distributed by their CD4 count results (cells per microliter).

First CD4 Test Result: First reported CD4 test result obtained within 12 months following initial HIV diagnosis.

Jurisdiction of Residence at AIDS Diagnosis: Jurisdiction of residence at time of initial AIDS diagnosis.

Jurisdiction of Residence at Diagnosis: Jurisdiction of residence at the later of time of initial HIV diagnosis or time of initial AIDS diagnosis.

Jurisdiction of Residence at HIV Diagnosis: Jurisdiction of residence at time of initial HIV diagnosis.

Mean Years from HIV Diagnosis (to AIDS Diagnosis): Mean number of years from initial HIV diagnosis to initial AIDS diagnosis for cases with a reported AIDS diagnosis.

Median: The measure of central location which divides a set of data into two equal parts.

Median Count (First CD4): Median CD4 count (cells per microliter) of the first CD4 test result reported within 12 months following initial HIV diagnosis.

Median Count (Recent CD4): Median CD4 count (cells per microliter) of the most recent CD4 test result reported in the 12 months prior to 12/31/2015.

Median Unsuppressed (Viral Load): Median unsuppressed viral load (copies per milliliter) among adult/adolescent living HIV cases with a most recent viral load test result reported in the 12 months prior to 12/31/2015 of 200 copies per milliliter or greater.

Percent Late HIV Diagnosis (for AIDS diagnoses): Percent of adult/adolescent reported AIDS diagnoses with an initial HIV diagnosis less than or equal to 12 months prior to their initial AIDS diagnosis.

Percent Late HIV Diagnosis (for HIV diagnoses): Percent of adult/adolescent reported HIV diagnoses with an initial AIDS diagnosis less than or equal to 12 months after their initial HIV diagnosis.

Percent Linked to Care: Percent of adult/adolescent reported HIV diagnoses with a CD4 or viral load test performed less than or equal to 3 months after their initial HIV diagnosis.

Percent Suppressed (Viral Load): Percent of adult/adolescent total living HIV cases with a most recent viral load reported in the 12 months prior to 12/31/2015 of less than 200 copies per milliliter.

Population Age 13+: Population age 13 years or older, estimate for 7/1/2014.

Rate: A proportion used to represent risk for disease within a given population. It is calculated by dividing the number of diagnoses by the number of persons at risk (population estimate).

Ratio (1 in X): Number of people for every 1 living HIV case in the population, or 1 living HIV case in every X number of people.

Recent CD4 Test Result: The most recent CD4 test result reported in the 12 months prior to 12/31/2015.

Recent Viral Load Test Result: The most recent viral load test result reported in the 12 months prior to 12/31/2015.

Suggested Citation: Maryland HIV/AIDS Quarterly Update, Second Quarter 2016. Baltimore, MD: Center for HIV Surveillance, Epidemiology and Evaluation, Infectious Disease Epidemiology and Outbreak Response Bureau, Prevention and Health Promotion Administration, Maryland Department of Health and Mental Hygiene. June 2016.

Section II – Adult/Adolescent Cases by Jurisdiction

Table 1 – Adult/Adolescent HIV Diagnoses during 1/1/2015-12/31/2015, First CD4 Test Result, Percent Linked to Care, and Percent Late Diagnosis, by Jurisdiction, Reported through 6/30/2016

JURISDICTION OF RESIDENCE AT HIV DIAGNOSIS	Population Age 13+	Adult/Adolescent Reported HIV Diagnoses							
		No.	% of Total	Rate	First CD4 Test Result			% Linked to Care	% Late HIV Diagnosis
					No. with Test	% with Test	Median Count		
Allegany	63,865	2	0.1%	3.1	***	***	***	***	***
Anne Arundel	468,309	66	4.9%	14.1	55	83.3%	329	81.8%	25.8%
Baltimore City	523,281	322	24.1%	61.5	250	77.6%	391	83.2%	21.4%
Baltimore	698,028	168	12.6%	24.1	138	82.1%	379	86.9%	25.0%
Calvert	75,818	3	0.2%	4.0	***	***	***	***	***
Caroline	26,987	3	0.2%	11.1	***	***	***	***	***
Carroll	142,505	6	0.4%	4.2	6	100.0%	141	100.0%	50.0%
Cecil	85,505	4	0.3%	4.7	***	***	***	***	***
Charles	128,189	30	2.2%	23.4	22	73.3%	433	80.0%	20.0%
Dorchester	27,470	10	0.7%	36.4	10	100.0%	167	90.0%	60.0%
Frederick	202,815	14	1.0%	6.9	12	85.7%	450	92.9%	14.3%
Garrett	25,684	2	0.1%	7.8	***	***	***	***	***
Harford	210,360	11	0.8%	5.2	8	72.7%	110	81.8%	45.5%
Howard	256,471	26	1.9%	10.1	21	80.8%	343	92.3%	30.8%
Kent	17,418	1	0.1%	5.7	***	***	***	***	***
Montgomery	855,169	205	15.4%	24.0	166	81.0%	352	83.4%	26.8%
Prince George's	753,706	391	29.3%	51.9	328	83.9%	375	84.9%	21.2%
Queen Anne's	41,277	1	0.1%	2.4	***	***	***	***	***
Saint Mary's	90,661	5	0.4%	5.5	***	***	***	***	***
Somerset	22,645	4	0.3%	17.7	***	***	***	***	***
Talbot	32,664	4	0.3%	12.2	***	***	***	***	***
Washington	126,009	10	0.7%	7.9	8	80.0%	682	90.0%	10.0%
Wicomico	85,525	15	1.1%	17.5	12	80.0%	256	73.3%	26.7%
Worcester	45,243	3	0.2%	6.6	***	***	***	***	***
Corrections	--	29	2.2%	--	24	82.8%	548	79.3%	10.3%
TOTAL	5,005,603	1,335	100.0%	26.7	1,087	81.4%	373	84.4%	23.7%

*** Data withheld due to low population counts and/or case counts

Adult/Adolescent Reported HIV Diagnoses: Reported HIV diagnoses, age 13 years or older at HIV diagnosis, with an initial HIV diagnosis during the specified year.

Jurisdiction of Residence at HIV Diagnosis: Jurisdiction of residence at time of initial HIV diagnosis.

Population Age 13+: Population age 13 years or older, estimate for 7/1/2014.

Rate: A proportion used to represent risk for disease within a given population. It is calculated by dividing the number of diagnoses by the number of persons at risk (population estimate).

First CD4 Test Result: First reported CD4 test result obtained within 12 months following initial HIV diagnosis.

Median Count (First CD4): Median CD4 count (cells per microliter) of the first CD4 test result reported within 12 months following initial HIV diagnosis.

Percent Linked to Care: Percent of adult/adolescent reported HIV diagnoses with a CD4 or viral load test performed less than or equal to 3 months after their initial HIV diagnosis.

Percent Late HIV Diagnosis (for HIV diagnoses): Percent of adult/adolescent reported HIV diagnoses with an initial AIDS diagnosis less than or equal to 12 months after their initial HIV diagnosis.

Table 2 – Adult/Adolescent AIDS Diagnoses during 1/1/2015-12/31/2015, Mean Years from HIV Diagnosis and Percent Late HIV Diagnosis, by Jurisdiction, Reported through 6/30/2016

JURISDICTION OF RESIDENCE AT AIDS DIAGNOSIS	Population Age 13+	Adult/Adolescent Reported AIDS Diagnoses				
		No.	% of Total	Rate	Mean Years from HIV Diagnosis	% Late HIV Diagnosis
Allegany	63,865	0	0.0%	0.0	--	--
Anne Arundel	468,309	32	5.0%	6.8	3.4	59.4%
Baltimore City	523,281	203	31.7%	38.8	5.1	38.9%
Baltimore	698,028	81	12.7%	11.6	4.8	46.9%
Calvert	75,818	3	0.5%	4.0	***	***
Caroline	26,987	3	0.5%	11.1	***	***
Carroll	142,505	5	0.8%	3.5	1.0	80.0%
Cecil	85,505	2	0.3%	2.3	***	***
Charles	128,189	10	1.6%	7.8	6.1	40.0%
Dorchester	27,470	9	1.4%	32.8	5.6	55.6%
Frederick	202,815	3	0.5%	1.5	***	***
Garrett	25,684	0	0.0%	0.0	--	--
Harford	210,360	10	1.6%	4.8	2.9	80.0%
Howard	256,471	14	2.2%	5.5	5.0	50.0%
Kent	17,418	2	0.3%	11.5	***	***
Montgomery	855,169	85	13.3%	9.9	2.9	68.2%
Prince George's	753,706	153	23.9%	20.3	3.4	55.6%
Queen Anne's	41,277	0	0.0%	0.0	--	--
Saint Mary's	90,661	4	0.6%	4.4	***	***
Somerset	22,645	2	0.3%	8.8	***	***
Talbot	32,664	0	0.0%	0.0	--	--
Washington	126,009	2	0.3%	1.6	***	***
Wicomico	85,525	7	1.1%	8.2	4.4	57.1%
Worcester	45,243	2	0.3%	4.4	***	***
Corrections	--	8	1.3%	--	8.0	37.5%
TOTAL	5,005,603	640	100.0%	12.8	4.2	51.6%

*** Data withheld due to low population counts and/or case counts

Adult/Adolescent Reported AIDS Diagnoses: Reported HIV diagnoses, age 13 years or older at HIV diagnosis, with an initial AIDS diagnosis during the specified year.

Jurisdiction of Residence at AIDS Diagnosis: Jurisdiction of residence at time of initial AIDS diagnosis.

Population Age 13+: Population age 13 years or older, estimate for 7/1/2014.

Rate: A proportion used to represent risk for disease within a given population. It is calculated by dividing the number of diagnoses by the number of persons at risk (population estimate).

Mean Years from HIV Diagnosis (to AIDS Diagnosis): Mean number of years from initial HIV diagnosis to initial AIDS diagnosis for cases with a reported AIDS diagnosis.

Percent Late HIV Diagnosis (for AIDS diagnoses): Percent of adult/adolescent reported AIDS diagnoses with an initial HIV diagnosis less than or equal to 12 months prior to their initial AIDS diagnosis.

Table 3 – Adult/Adolescent HIV Cases Alive on 12/31/2015, by Jurisdiction, Reported through 6/30/2016

JURISDICTION OF RESIDENCE AT DIAGNOSIS	Population Age 13+	Adult/Adolescent Living HIV Cases without AIDS			Adult/Adolescent Living HIV Cases with AIDS			Adult/Adolescent Total Living HIV Cases			
		No.	% of Total	Rate	No.	% of Total	Rate	No.	% of Total	Rate	Ratio (1 in X)
Allegany	63,865	42	0.3%	65.8	32	0.2%	50.1	74	0.2%	115.9	863
Anne Arundel	468,309	565	3.8%	120.6	671	3.9%	143.3	1,236	3.9%	263.9	378
Baltimore City	523,281	5,382	36.2%	1,028.5	6,591	38.7%	1,259.6	11,973	37.6%	2,288.1	43
Baltimore	698,028	1,553	10.5%	222.5	1,725	10.1%	247.1	3,278	10.3%	469.6	212
Calvert	75,818	47	0.3%	62.0	50	0.3%	65.9	97	0.3%	127.9	781
Caroline	26,987	30	0.2%	111.2	36	0.2%	133.4	66	0.2%	244.6	408
Carroll	142,505	61	0.4%	42.8	72	0.4%	50.5	133	0.4%	93.3	1,071
Cecil	85,505	51	0.3%	59.6	60	0.4%	70.2	111	0.3%	129.8	770
Charles	128,189	252	1.7%	196.6	197	1.2%	153.7	449	1.4%	350.3	285
Dorchester	27,470	41	0.3%	149.3	83	0.5%	302.1	124	0.4%	451.4	221
Frederick	202,815	173	1.2%	85.3	157	0.9%	77.4	330	1.0%	162.7	614
Garrett	25,684	6	0.0%	23.4	4	0.0%	15.6	10	0.0%	38.9	2,568
Harford	210,360	190	1.3%	90.3	236	1.4%	112.2	426	1.3%	202.5	493
Howard	256,471	251	1.7%	97.9	254	1.5%	99.0	505	1.6%	196.9	507
Kent	17,418	16	0.1%	91.9	19	0.1%	109.1	35	0.1%	200.9	497
Montgomery	855,169	1,892	12.7%	221.2	2,054	12.1%	240.2	3,946	12.4%	461.4	216
Prince George's	753,706	3,361	22.6%	445.9	3,479	20.4%	461.6	6,840	21.5%	907.5	110
Queen Anne's	41,277	14	0.1%	33.9	33	0.2%	79.9	47	0.1%	113.9	878
Saint Mary's	90,661	63	0.4%	69.5	65	0.4%	71.7	128	0.4%	141.2	708
Somerset	22,645	25	0.2%	110.4	31	0.2%	136.9	56	0.2%	247.3	404
Talbot	32,664	28	0.2%	85.7	32	0.2%	98.0	60	0.2%	183.7	544
Washington	126,009	169	1.1%	134.1	129	0.8%	102.4	298	0.9%	236.5	422
Wicomico	85,525	102	0.7%	119.3	109	0.6%	127.4	211	0.7%	246.7	405
Worcester	45,243	29	0.2%	64.1	47	0.3%	103.9	76	0.2%	168.0	595
Corrections	--	518	3.5%	--	848	5.0%	--	1,366	4.3%	--	--
TOTAL	5,005,603	14,861	100.0%	296.9	17,014	100.0%	339.9	31,875	100.0%	636.8	157

Jurisdiction of Residence at Diagnosis: Jurisdiction of residence at later of time of initial HIV diagnosis or time of initial AIDS diagnosis.

Population Age 13+: Population greater than or equal to 13 years old, estimate for 7/1/2014.

Adult/Adolescent Living HIV Cases without AIDS: Reported HIV diagnoses, age 13 years or older at HIV diagnosis, without an AIDS diagnosis, and not reported to have died as of 12/31/2015.

Adult/Adolescent Living HIV Cases with AIDS: Reported HIV diagnoses, age 13 years or older at HIV diagnosis, with an AIDS diagnosis, and not reported to have died as of 12/31/2015.

Adult/Adolescent Total Living HIV Cases: Reported HIV diagnoses, age 13 years or older at HIV diagnosis, with or without an AIDS diagnosis, and not reported to have died as of 12/31/2015.

Rate: A proportion used to represent risk for disease within a given population. It is calculated by dividing the number of diagnoses by the number of persons at risk (population estimate).

Ratio (1 in X): Number of people for every 1 living HIV case in the population, or 1 living HIV case in every X number of people.

Table 4 – CD4 Test Results for Adult/Adolescent HIV Cases Alive on 12/31/2015, Reported through 6/30/2016

JURISDICTION OF RESIDENCE AT DIAGNOSIS	Adult/Adolescent Total Living HIV Cases								
	No.	Recent CD4 Test Result						350-499	500+
		No. with Test	% with Test	Median Count	<200	200-349			
Allegany	74	50	67.6%	652	4.0%	8.0%	20.0%	68.0%	
Anne Arundel	1,236	730	59.1%	548	12.5%	15.1%	17.7%	54.8%	
Baltimore City	11,973	6,782	56.6%	528	14.1%	15.2%	17.5%	53.3%	
Baltimore	3,278	1,892	57.7%	556	12.2%	13.6%	17.2%	57.0%	
Calvert	97	67	69.1%	642	9.0%	11.9%	11.9%	67.2%	
Caroline	66	29	43.9%	476	17.2%	10.3%	27.6%	44.8%	
Carroll	133	64	48.1%	595	20.3%	10.9%	7.8%	60.9%	
Cecil	111	49	44.1%	546	6.1%	22.4%	16.3%	55.1%	
Charles	449	269	59.9%	610	9.3%	12.6%	15.2%	62.8%	
Dorchester	124	83	66.9%	509	10.8%	18.1%	19.3%	51.8%	
Frederick	330	186	56.4%	569	8.6%	11.3%	19.9%	60.2%	
Garrett	10	6	60.0%	679	16.7%	0.0%	0.0%	83.3%	
Harford	426	256	60.1%	553	14.1%	15.6%	16.0%	54.3%	
Howard	505	277	54.9%	563	12.3%	16.6%	14.8%	56.3%	
Kent	35	21	60.0%	680	9.5%	28.6%	4.8%	57.1%	
Montgomery	3,946	2,073	52.5%	542	10.0%	14.1%	20.4%	55.5%	
Prince George's	6,840	3,792	55.4%	540	11.6%	14.4%	18.9%	55.1%	
Queen Anne's	47	27	57.4%	605	14.8%	18.5%	3.7%	63.0%	
Saint Mary's	128	84	65.6%	461	11.9%	14.3%	27.4%	46.4%	
Somerset	56	41	73.2%	553	12.2%	12.2%	14.6%	61.0%	
Talbot	60	30	50.0%	394	23.3%	23.3%	10.0%	43.3%	
Washington	298	160	53.7%	718	7.5%	7.5%	13.8%	71.3%	
Wicomico	211	126	59.7%	520	18.3%	11.9%	17.5%	52.4%	
Worcester	76	50	65.8%	577	10.0%	10.0%	18.0%	62.0%	
Corrections	1,366	770	56.4%	475	18.2%	17.1%	17.9%	46.8%	
TOTAL	31,875	17,914	56.2%	539	12.7%	14.6%	18.0%	54.7%	

Adult/Adolescent Total Living HIV Cases: Reported HIV diagnoses, age 13 years or older at HIV diagnosis, with or without an AIDS diagnosis, and not reported to have died as of 12/31/2015.

Recent CD4 Test Result: The most recent CD4 test result reported in the 12 months prior to 12/31/2015.

Jurisdiction of Residence at Diagnosis: Jurisdiction of residence at later of time of initial HIV diagnosis or time of initial AIDS diagnosis.

Median Count (Recent CD4): Median CD4 count (cells per microliter) of the most recent CD4 test result reported in the 12 months prior to 12/31/2015.

CD4 Result Distribution (<200, 200-349, 350-499, 500+): Percent of cases with a CD4 test distributed by their CD4 count results (cells per microliter).

Table 5 – Viral Load Test Results for Adult/Adolescent HIV Cases Alive on 12/31/2015, by Jurisdiction, Reported through 6/30/2016

JURISDICTION OF RESIDENCE AT DIAGNOSIS	Adult/Adolescent Total Living HIV Cases				
	No.	Recent Viral Load Test Result			
		No. with Test	% with Test	% Suppressed	Median Unsuppressed
Allegany	74	51	68.9%	92.2%	22,422
Anne Arundel	1,236	682	55.2%	75.5%	12,076
Baltimore City	11,973	6,197	51.8%	70.7%	11,224
Baltimore	3,278	1,756	53.6%	76.9%	11,700
Calvert	97	69	71.1%	87.0%	31,664
Caroline	66	29	43.9%	79.3%	5,159
Carroll	133	67	50.4%	85.1%	45,252
Cecil	111	44	39.6%	81.8%	1,170
Charles	449	274	61.0%	76.6%	14,661
Dorchester	124	81	65.3%	84.0%	4,850
Frederick	330	193	58.5%	83.9%	4,427
Garrett	10	6	60.0%	83.3%	565
Harford	426	237	55.6%	78.5%	15,780
Howard	505	281	55.6%	78.3%	30,990
Kent	35	21	60.0%	85.7%	1,506
Montgomery	3,946	2,061	52.2%	85.2%	11,388
Prince George's	6,840	3,782	55.3%	79.0%	15,790
Queen Anne's	47	26	55.3%	88.5%	101,486
Saint Mary's	128	80	62.5%	78.8%	23,689
Somerset	56	38	67.9%	81.6%	10,410
Talbot	60	35	58.3%	77.1%	45,988
Washington	298	158	53.0%	81.0%	964
Wicomico	211	121	57.3%	72.7%	34,050
Worcester	76	50	65.8%	86.0%	9,680
Corrections	1,366	598	43.8%	62.0%	15,230
TOTAL	31,875	16,937	53.1%	75.9%	12,679

Adult/Adolescent Total Living HIV Cases: Reported HIV diagnoses, age 13 years or older at HIV diagnosis, with or without an AIDS diagnosis, and not reported to have died as of 12/31/2015.

Jurisdiction of Residence at Diagnosis: Jurisdiction of residence at later of time of initial HIV diagnosis or time of initial AIDS diagnosis.

Recent Viral Load Test Result: The most recent viral load test result reported in the 12 months prior to 12/31/2015.

Percent Suppressed (Viral Load): Percent of adult/adolescent total living HIV cases with a most recent viral load reported in the 12 months prior to 12/31/2015 of less than 200 copies per milliliter.

Median Unsuppressed (Viral Load): Median unsuppressed viral load (copies per milliliter) among adult/adolescent living HIV cases with a most recent viral load test result reported in the 12 months prior to 12/31/2015 of 200 copies per milliliter or greater.