

Weekly Influenza Surveillance Report

Maryland Department of Health and Mental Hygiene | Infectious Disease and Environmental Health Administration
Office of Infectious Disease Epidemiology and Outbreak Response

SYNOPSIS

During the week of April 17 – 23 (week 16) influenza activity in Maryland was “sporadic”. For the first time in quite a while, none of the surveillance regions had substantial influenza activity. However, the State Laboratories Administration continues to report PCR-positive specimens for influenza types A and B. So the virus is still circulating in Maryland, but it is doing so at low levels. Two hospitalizations associated with influenza were reported during week 16. No outbreaks of respiratory disease were reported during week 16. These weekly influenza reports will continue until week 20 (May 21). After that date, surveillance will continue, but the reports will be monthly.

PLEASE NOTE: Influenza is not a reportable condition in Maryland. As a result, we rely on select sources of information such as some (sentinel) clinical labs and physician offices, and the public. Because these sources cannot report all cases in the state, the counts contained in this summary do not represent the true number of cases of influenza in Maryland. They do provide valuable information about trends. All data are preliminary and subject to change.

INFLUENZA-LIKE ILLNESS SURVEILLANCE (ILINet)

During week 16, seven sentinel providers reported 127 (2.3%) of 5,534 visits to their practices were for ILI. This is below the state baseline of 5.6%.

This same week last season, when influenza activity peaked late in October of 2009 and was on the decline by December, the proportion of visits for ILI was 1.2%.

For more information on the US Outpatient Influenza-like Illness Reporting Network (ILINet), please visit our website: <http://dhmh.maryland.gov/fluwatch> and click on “ILINet Sentinel Providers”.




Figure 1. Proportion of visits for ILI to ILINet sentinel providers, 2010-11 influenza season

CLINICAL LAB REPORTS OF RAPID FLU TESTING

During week 16, nineteen sentinel clinical laboratories reported 30 (6.2%) of 487 rapid influenza tests as positive: 8 were positive for type A, and 22 were positive for type B influenza. This proportion of positive tests is higher than the proportion reported at this time last season (2009-2010), which was 1.3%, and higher compared to the proportion observed during the same week of the 2008-2009 season.

While not as accurate as PCR tests, rapid influenza tests become more accurate as the flu season progresses and as influenza becomes more prevalent in the community, giving insight into the activity of influenza.




Figure 2. Proportion of positive rapid tests reported by sentinel clinical laboratories, 2010-11 influenza season

Type of Positives	Number (%)
Type A	4,837 (79%)
Type B	1,274 (21%)
Positive, but not typed	0
Total Positive	6,111 (100%)

Table 1. Number of positive rapid influenza tests, by type, reported by collaborating clinical laboratories 2010-11 season to date

GET VACCINATED!
Go to
<http://dhmh.maryland.gov/swineflu/getVaccinated.html>
and find your local health department for more information.

MARYLAND RESIDENT INFLUENZA TRACKING SURVEY (MRITS)

During week 16, a total of 497 (33.4% of total) participants in the MRITS responded to the weekly survey. Of those who responded, 10 (2.0%) reported flu-like illness. This proportion is higher than this same week last season, when 1.7% of respondents reported flu-like illness.

We are always looking for more participants for the MRITS. If you know someone who would like to participate, please direct them to our website: <http://dhmh.maryland.gov/flusurvey>.




Figure 3. Proportion reporting ILI to the MRITS by week, 2010-11 influenza season

DHMH LABORATORIES ADMINISTRATION REPORTS

During week 16, the DHMH Laboratories Administration performed a total of 34 PCR tests for influenza. Thirteen were positive for influenza: 1 was type A (H1N1), 2 were type A (H3), and 10 were type B.

The table to the right shows the breakdown of positive tests by influenza strain for the 2010-11 influenza season to date.

More information on the valuable work done by the DHMH Laboratories Administration is available at <http://dhmh.maryland.gov/labs>.

Influenza Type	No. (%)
Type A	
H1	716 (45.6%)
H3	722 (46.0%)
Unsubtyped	0 (0%)
Type B	132 (8.4%)
TOTAL	1,570 (100%)

Table 1. Number of respiratory samples positive for influenza by PCR reported by the DHMH Labs Administration, 2010-11 influenza season

EIP INFLUENZA HOSPITALIZATION SURVEILLANCE

During week 16, two (2) hospitalizations associated with influenza were reported to the Emerging Infections Program (EIP) by 39 hospitals. To date, there have been 1,211 hospitalizations.

During the same week last season, 2 hospitalizations were reported, with a total of 1,454 at that point in the season. For the entire season (2009-10), 1,458 hospitalizations were reported.

To be a confirmed hospitalization associated with influenza, the person must be hospitalized and have a positive influenza test of any kind (rapid test, PCR, culture).




Figure 4. Number of hospitalizations associated with influenza, by age group and week, reported to the Emerging Infections Program, 2010-11 influenza season

REPORTS OF OUTBREAKS IN INSTITUTIONAL SETTINGS

During week 16, no outbreaks of respiratory disease were reported to DHMH. This keeps the season’s total at 78 reported outbreaks. Last season, a total of 208 outbreaks of respiratory illness were reported. Of those, 33 were confirmed as influenza outbreaks.

An outbreak of ILI is re-classified as an outbreak of influenza if there is laboratory evidence of influenza virus present in the samples collected from case-patients during the outbreak.




Figure 5. Number of outbreaks reported by week and by type during the 2010-11 influenza season.

EMERGENCY DEPARTMENT ILI REPORTS (ESSENCE)

During week 16, a total of 41,997 visits to emergency departments for all reasons were reported to the Office of Preparedness and Response through the ESSENCE system. Of those visits, 400 (0.95%) were for influenza-like illness. This proportion is slightly lower than those observed over the prior two influenza seasons and slightly lower than the previous week.

For more information on ESSENCE, please visit the Office of Preparedness and Response’s web site at: <http://bioterrorism.dhmh.state.md.us>.




Figure 6. Proportion of visits to emergency departments for ILI by week reported through ESSENCE, 2010-11 influenza season.

GOOGLE FLU TRENDS

According to Google, influenza activity in Maryland is currently “MINIMAL”. What does this mean? From the [Google Flu Trends Website](http://www.google.com/flutrends/): “We have found a close relationship between how many people search for flu-related topics and how many people actually have flu symptoms. Of course, not every person who searches for ‘flu’ is actually sick, but a pattern emerges when all the flu-related search queries are added together. We compared our query counts with traditional flu surveillance systems and found that many search queries tend to be popular exactly when flu season is happening. By counting how often we see these search queries, we can estimate how much flu is circulating in different countries and regions around the world.”




Figure 7 – According to Google Flu Trends, influenza activity in Maryland is currently “minimal”. At this time last year, during the 2009 H1N1 influenza pandemic, influenza activity in Maryland was “minimal” to “low”.

