



Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to DHMH for the week ending April 9, 2016

Prepared by the Infectious Disease Epidemiology and Outbreak Response Bureau
Prevention and Health Promotion Administration
Maryland Department of Health and Mental Hygiene

The data presented in this document are provisional and subject to change as additional reports are received.

SUMMARY

During the week ending April 9, 2016, influenza-like illness (ILI) intensity in Maryland was **MINIMAL** and there was **REGIONAL** geographic spread. The proportion of outpatient visits for ILI reported by Sentinel Providers increased, while the proportion reported by Maryland Emergency Departments dropped for the third consecutive week. The proportion of MRITS respondents reporting ILI increased. The proportion of specimens testing positive for influenza at clinical laboratories dropped, but remained high. Fewer specimens tested positive for influenza at the DHMH lab, but the percent of specimens testing positive increased slightly. There were 47 influenza-associated hospitalizations reported. Two respiratory outbreaks were reported. Nationally, influenza activity continued to decline, but remained elevated.


[Click here to visit our influenza surveillance web page](#)

ILI Intensity Levels
✓ Minimal
Low
Moderate
High

Influenza Geographic Spread
No Activity
Sporadic
Local
✓ Regional
Widespread

ILINet Sentinel Providers


Eighteen sentinel providers reported a total of 4,421 visits this week. Of those, 81 (1.8%) were visits for ILI. This is **at** the baseline level for Maryland.



ILI Visits To Sentinel Providers By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	30 (37%)	19 (21%)	1004 (24%)
Age 5-24	20 (25%)	37 (40%)	1492 (36%)
Age 25-49	20 (25%)	18 (20%)	917 (22%)
Age 50-64	9 (11%)	13 (14%)	494 (12%)
Age ≥ 65	2 (2%)	5 (5%)	204 (5%)
Total	81 (100%)	92 (100%)	4111 (100%)

Visits to Emergency Departments for ILI

Emergency Departments in Maryland reported a total of 47,699 visits this week through the [ESSENCE surveillance system](#). Of those, 1,018 (2.1%) were visits for ILI.



ILI Visits To Emergency Departments By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	213 (21%)	302 (24%)	8291 (28%)
Age 5-24	299 (29%)	319 (26%)	8618 (30%)
Age 25-49	290 (28%)	375 (30%)	7555 (26%)
Age 50-64	126 (12%)	160 (13%)	2958 (10%)
Age ≥ 65	90 (9%)	94 (8%)	1713 (6%)
Unknown	--	--	--
Total	1018 (100%)	1250 (100%)	29135(100%)

Neighboring states' influenza information:


- Delaware <http://dhss.delaware.gov/dph/epi/influenzahome.html>
- District of Columbia <http://doh.dc.gov/service/influenza>
- Pennsylvania [http://www.portal.state.pa.us/portal/server.pt/community/influenza_\(flu\)/14161](http://www.portal.state.pa.us/portal/server.pt/community/influenza_(flu)/14161)
- Virginia <http://www.vdh.state.va.us/Epidemiology/flu/>
- West Virginia <http://dhhr.wv.gov/oeps/disease/flu/Pages/fluSurveillance.aspx>

Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to DHMH for the week ending April 9, 2016

Community-based Influenza Surveillance (MRITS)


MRITS is the Maryland Resident Influenza Tracking System, a weekly survey for influenza-like illness (ILI). A total of 594 residents responded to the [MRITS survey](#) this week. Of those, 10 (1.7%) reported having ILI and missing a cumulative 19 days of regular daily activities.




MRITS Respondents Reporting ILI By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	1 (10%)	--	14 (6%)
Age 5-24	5 (50%)	1 (14%)	53 (22%)
Age 25-49	2 (20%)	3 (43%)	67 (28%)
Age 50-64	1 (10%)	2 (29%)	77 (32%)
Age ≥ 65	1 (10%)	1 (14%)	31 (13%)
Total	10 (100%)	7 (100%)	242 (100%)

Clinical Laboratory Influenza Testing

Forty-three clinical laboratories reported performing 2,825 influenza diagnostic tests, mostly rapid influenza diagnostic tests (RIDTs). Of those, 651 (23.0%) were positive for influenza. Of those testing positive, 324 (49.8%) were influenza type A and 327 (50.2%) were influenza type B. The [reliability of RIDTs](#) depends largely on the conditions under which they are used. False-positive (and true-negative) results are more likely to occur when the disease prevalence in the community is low, which is generally at the beginning and end of the influenza season and during the summer.




Positive Rapid Flu Tests by Type	This Week Number (%)	Last Week Number (%)	Season Number (%)
Type A	324 (50%)	563 (59%)	6141 (70%)
Type B	327 (50%)	398 (41%)	2668 (30%)
Total	651 (100%)	961 (100%)	8809 (100%)



State Laboratories Administration Influenza Testing

The DHMH Laboratories Administration performed a total of 203 PCR tests for influenza and 105 (51.7%) specimens tested positive for influenza. Of those testing positive, 75 (71.4%) were type A (H1), 17 (16.2%) were type B (Yamagata), and 13 (12.4%) were type B (Victoria). PCR testing is more reliable than RIDT. The DHMH testing identifies subtypes of influenza A and lineages of influenza B, information that is not available from the RIDT results. The lab performs genetic sequencing for a subset of the specimens they receive and can detect genes associated with antiviral resistance within those sequences.



Positive PCR Tests by Type (Subtype)	This Week Number (%)	Last Week Number (%)	Season Number (%)
Type A (H1)	75 (71%)	87 (70%)	1053 (81%)
Type A (H3)	--	1 (1%)	36 (3%)
Type B (Victoria)	13 (12%)	14 (11%)	92 (7%)
Type B (Yamagata)	17 (16%)	22 (18%)	124 (9%)
Dual - Type A(H3) / B (Yamagata)	--	--	1 (<1%)
Total	105 (100%)	124 (100%)	1306 (100%)

Gene Mutation Associated w/ Antiviral Resistance	This Week Number (%)	Last Week Number (%)	Season Number (%)
Detected	--	--	1 (<1%)
Not Detected	--	--	263 (100%)
Total	--	--	264 (100%)

Where to get an influenza vaccination


Interested in getting a flu vaccine for the 2015-16 influenza season? Go to <http://dhhm.maryland.gov/flum/d/Pages/getvaccinated.aspx> and click on your county/city of residence. You will be redirected to your local health department website for local information on where to get your flu vaccine.

Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to DHMH for the week ending April 9, 2016

Influenza-associated Hospitalizations


There were 47 influenza-associated hospitalizations reported to DHMH this week. (A person with an overnight hospital stay along with a positive influenza test of any kind, e.g. RIDT or PCR, is considered an “influenza-associated hospitalization” for purposes of influenza surveillance.)



Influenza-Associated Hospitalizations by Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	4 (9%)	13 (11%)	132 (10%)
Age 5-17	2 (4%)	5 (4%)	78 (6%)
Age 18-24	1 (2%)	5 (4%)	41 (3%)
Age 25-49	10 (21%)	26 (23%)	283 (22%)
Age 50-64	13 (28%)	22 (19%)	360 (28%)
Age ≥ 65	17 (36%)	44 (38%)	391 (30%)
Total	47 (100%)	115 (100%)	1285 (100%)

Outbreaks of Respiratory Disease

There were 2 influenza outbreaks reported to DHMH this week. (Disease outbreaks of any kind are reportable in Maryland. Respiratory outbreaks may be reclassified once a causative agent is detected, e.g. from ILI to influenza.)



Respiratory Outbreaks by Type	This Week Number (%)	Last Week Number (%)	Season Number (%)
Influenza	1 (50%)	1 (25%)	12 (39%)
Influenza-like Illness	1 (50%)	1 (25%)	7 (23%)
Pneumonia	--	2 (50%)	12 (39%)
Other Respiratory	--	--	--
Total	2 (100%)	4 (100%)	31 (100%)

National Influenza Surveillance (CDC)

During week 14 (April 3-9, 2016), influenza activity decreased, but remained elevated in the United States.

- Viral Surveillance:** The most frequently identified influenza virus type reported by public health laboratories during week 14 was influenza A, with influenza A (H1N1)pdm09 viruses predominating. The percentage of respiratory specimens testing positive for influenza in clinical laboratories decreased.
- Pneumonia and Influenza Mortality:** The proportion of deaths attributed to pneumonia and influenza (P&I) was below the system-specific epidemic threshold in the NCHS Mortality Surveillance System and above the system-specific epidemic threshold in the 122 Cities Mortality Reporting System.
- Influenza-associated Pediatric Deaths:** Ten influenza-associated pediatric deaths were reported.
- Influenza-associated Hospitalizations:** A cumulative rate for the season of 26.6 laboratory-confirmed influenza-associated hospitalizations per 100,000 population was reported.
- Outpatient Illness Surveillance:** The proportion of outpatient visits for influenza-like illness (ILI) was 2.1%, which is at the national baseline of 2.1%. Six of 10 regions reported ILI at or above region-specific baseline levels. One state experienced high ILI activity; Puerto Rico and one state experienced moderate ILI activity; 11 states experienced low ILI activity; New York City and 36 states experienced minimal ILI activity; and the District of Columbia and one state had insufficient data.
- Geographic Spread of Influenza:** The geographic spread of influenza in Puerto Rico and 18 states was reported as widespread; Guam and 19 states reported regional activity; the District of Columbia and ten states reported local activity; and the U.S. Virgin Islands and three states reported sporadic activity.

