



# Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to DHMH for the week ending March 26, 2016

Prepared by the Infectious Disease Epidemiology and Outbreak Response Bureau  
Prevention and Health Promotion Administration  
Maryland Department of Health and Mental Hygiene

*The data presented in this document are provisional and subject to change as additional reports are received.*

## SUMMARY

During the week ending March 26, 2016, influenza-like illness (ILI) intensity in Maryland was **MINIMAL** and there was **WIDESPREAD** geographic spread. The proportion of outpatient visits for ILI reported by both Sentinel Providers and Maryland Emergency Departments dropped. The proportion of MRITS respondents reporting ILI decreased. The proportion of specimens testing positive for influenza at clinical laboratories experienced a slight decrease. There was a large decrease in the number of specimens testing positive and in the percent positivity for influenza at the DHMH lab. There were 94 influenza-associated hospitalizations reported. One influenza outbreak was reported. Nationally, influenza activity is declining.


[Click here to visit our influenza surveillance web page](#)

ILI Intensity Levels
✓ Minimal
Low
Moderate
High

Influenza Geographic Spread
No Activity
Sporadic
Local
Regional
✓ Widespread

## ILINet Sentinel Providers


Twenty-two sentinel providers reported a total of 7,198 visits this week. Of those, 94 (1.3%) were visits for ILI. This is below the Maryland baseline of 1.8%.



ILI Visits To Sentinel Providers By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	27 (29%)	41 (30%)	942 (25%)
Age 5-24	38 (40%)	56 (41%)	1404 (37%)
Age 25-49	19 (20%)	23 (17%)	840 (22%)
Age 50-64	8 (9%)	14 (10%)	446 (12%)
Age ≥ 65	2 (2%)	3 (2%)	187 (5%)
Total	94 (100%)	137 (100%)	3819 (100%)

## Visits to Emergency Departments for ILI

Emergency Departments in Maryland reported a total of 49,463 visits this week through the [ESSENCE surveillance system](#). Of those, 1,647 (3.3%) were visits for ILI.



ILI Visits To Emergency Departments By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	389 (24%)	424 (22%)	7776 (29%)
Age 5-24	494 (30%)	620 (32%)	8000 (30%)
Age 25-49	476 (29%)	561 (29%)	6890 (26%)
Age 50-64	181 (11%)	216 (11%)	2672 (10%)
Age ≥ 65	107 (6%)	132 (7%)	1529 (6%)
Unknown	--	--	--
Total	1647 (100%)	1953 (100%)	26867(100%)

## Neighboring states' influenza information:


- Delaware <http://dhss.delaware.gov/dph/epi/influenzahome.html>
- District of Columbia <http://doh.dc.gov/service/influenza>
- Pennsylvania [http://www.portal.state.pa.us/portal/server.pt/community/influenza\\_\(flu\)/14161](http://www.portal.state.pa.us/portal/server.pt/community/influenza_(flu)/14161)
- Virginia <http://www.vdh.state.va.us/Epidemiology/flu/>
- West Virginia <http://dhhr.wv.gov/oeps/disease/flu/Pages/fluSurveillance.aspx>

# Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to DHMH for the week ending March 26, 2016

## Community-based Influenza Surveillance (MRITS)


MRITS is the Maryland Resident Influenza Tracking System, a weekly survey for influenza-like illness (ILI). A total of 603 residents responded to the [MRITS survey](#) this week. Of those, 9 (1.5%) reported having ILI and missing a cumulative 17 days of regular daily activities.




MRITS Respondents Reporting ILI By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	--	1 (7%)	13 (6%)
Age 5-24	3 (33%)	1 (7%)	47 (21%)
Age 25-49	5 (56%)	5 (36%)	62 (28%)
Age 50-64	1 (11%)	3 (21%)	74 (33%)
Age ≥ 65	--	4 (29%)	29 (13%)
<b>Total</b>	<b>9 (100%)</b>	<b>14 (100%)</b>	<b>225 (100%)</b>

## Clinical Laboratory Influenza Testing

Forty clinical laboratories reported performing 4,252 influenza diagnostic tests, mostly rapid influenza diagnostic tests (RIDTs). Of those, 1,161 (27.3%) were positive for influenza. Of those testing positive, 747 (64.3%) were influenza type A and 414 (35.7%) were influenza type B. The [reliability of RIDTs](#) depends largely on the conditions under which they are used. False-positive (and true-negative) results are more likely to occur when the disease prevalence in the community is low, which is generally at the beginning and end of the influenza season and during the summer.




Positive Rapid Flu Tests by Type	This Week Number (%)	Last Week Number (%)	Season Number (%)
Type A	747 (64%)	972 (70%)	5246 (73%)
Type B	414 (36%)	426 (30%)	1940 (27%)
<b>Total</b>	<b>1161 (100%)</b>	<b>1398 (100%)</b>	<b>7186 (100%)</b>



## State Laboratories Administration Influenza Testing

The DHMH Laboratories Administration performed a total of 339 PCR tests for influenza and 163 (48.1%) specimens tested positive for influenza. Of those testing positive, 111 (68.1%) were type A (H1), 3 (1.8%) were type A (H3), 26 (16.0%) were type B (Yamagata), 22 (13.5%) were type B (Victoria), and there was 1 (0.6%) dual infection with type A (H3) and B (Yamagata). PCR testing is more reliable than RIDT. The DHMH testing identifies subtypes of influenza A and lineages of influenza B, information that is not available from the RIDT results. The lab performs genetic sequencing for a subset of the specimens they receive and can detect genes associated with antiviral resistance within those sequences.



Positive PCR Tests by Type (Subtype)	This Week Number (%)	Last Week Number (%)	Season Number (%)
Type A (H1)	111 (68%)	175 (81%)	891 (83%)
Type A (H3)	3 (2%)	8 (4%)	36 (3%)
Type B (Victoria)	22 (13%)	16 (7%)	65 (6%)
Type B (Yamagata)	26 (16%)	16 (7%)	86 (8%)
Dual - Type A(H3) / B (Yamagata)	1 (1%)	--	1 (<1%)
<b>Total</b>	<b>163 (100%)</b>	<b>215 (100%)</b>	<b>1077 (100%)</b>

Gene Mutation Associated w/Antiviral Resistance	This Week Number (%)	Last Week Number (%)	Season Number (%)
Detected	--	--	1 (<1%)
Not Detected	--	--	263 (100%)
<b>Total</b>	<b>--</b>	<b>--</b>	<b>264 (100%)</b>

## Where to get an influenza vaccination


Interested in getting a flu vaccine for the 2015-16 influenza season? Go to <http://dhhm.maryland.gov/flumd/Pages/getvaccinated.aspx> and click on your county/city of residence. You will be redirected to your local health department website for local information on where to get your flu vaccine.

# Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to DHMH for the week ending March 26, 2016

## Influenza-associated Hospitalizations


There were 94 influenza-associated hospitalizations reported to DHMH this week. (A person with an overnight hospital stay along with a positive influenza test of any kind, e.g. RIDT or PCR, is considered an “influenza-associated hospitalization” for purposes of influenza surveillance.)



Influenza-Associated Hospitalizations by Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	10 (11%)	19 (12%)	105 (10%)
Age 5-17	8 (9%)	7 (4%)	65 (6%)
Age 18-24	1 (1%)	7 (4%)	30 (3%)
Age 25-49	14 (15%)	38 (23%)	223 (22%)
Age 50-64	30 (32%)	45 (28%)	293 (29%)
Age ≥ 65	31 (33%)	47 (29%)	297 (29%)
<b>Total</b>	<b>94 (100%)</b>	<b>163 (100%)</b>	<b>1013 (100%)</b>

## Outbreaks of Respiratory Disease

There was 1 influenza outbreak reported to DHMH this week. (Disease outbreaks of any kind are reportable in Maryland. Respiratory outbreaks may be reclassified once a causative agent is detected, e.g. from ILI to influenza.)



Respiratory Outbreaks by Type	This Week Number (%)	Last Week Number (%)	Season Number (%)
Influenza	1 (100%)	3 (75%)	10 (40%)
Influenza-like illness	--	1 (25%)	5 (20%)
Pneumonia	--	--	10 (40%)
Other Respiratory	--	--	--
<b>Total</b>	<b>1 (100%)</b>	<b>4 (100%)</b>	<b>25 (100%)</b>

## National Influenza Surveillance (CDC)

During week 12 (March 20-26, 2016), influenza activity decreased slightly, but remained elevated in the United States.

- Viral Surveillance:** The most frequently identified influenza virus type reported by public health laboratories during week 12 was influenza A, with influenza A (H1N1)pdm09 viruses predominating. The percentage of respiratory specimens testing positive for influenza in clinical laboratories decreased.
- Pneumonia and Influenza Mortality:** The proportion of deaths attributed to pneumonia and influenza (P&I) was below the system-specific epidemic threshold in the NCHS Mortality Surveillance System and above the system-specific epidemic threshold in the 122 Cities Mortality Reporting System.
- Influenza-associated Pediatric Deaths:** Three influenza-associated pediatric deaths were reported.
- Influenza-associated Hospitalizations:** A cumulative rate for the season of 21.4 laboratory-confirmed influenza-associated hospitalizations per 100,000 population was reported.
- Outpatient Illness Surveillance:** The proportion of outpatient visits for influenza-like illness (ILI) was 2.9%, which is above the national baseline of 2.1%. Nine of 10 regions reported ILI at or above region-specific baseline levels. Puerto Rico and two states experienced high ILI activity; New York City and seven states experienced moderate ILI activity; 15 states experienced low ILI activity; 26 states experienced minimal ILI activity; and the District of Columbia had insufficient data.
- Geographic Spread of Influenza:** The geographic spread of influenza in Guam, Puerto Rico and 29 states was reported as widespread; 18 states reported regional activity; the District of Columbia and two states reported local activity; one state reported sporadic activity; and the U.S. Virgin Islands did not report.

